

The effects of chronic carbon monoxide treatment on diet-induced obesity and *Rev-erb-alpha* target gene expression in adipose tissue

Tim Brodsky, Ryan McLoughlin, Ben Sell, and Thomas H. Reynolds IV*

Department of Health and Exercise Sciences, Skidmore College, USA

Abstract

The present study determined if carbon monoxide (CO) treatment reversed diet-induced obesity and insulin resistance. Because CO is a potential ligand for the nuclear receptor *Rev-erb-α*, we also determined if CO treatment altered the mRNA expression of *Rev-erb-α* targets. Mice were fed a low fat diet (LFD) or a high fat diet (HFD) for 150 days. Following 110 days of the dietary intervention the HFD-fed mice were assigned to a CO inhalation group or a control group. The HFD-CO group was exposed to 250 ppm CO for one hour per day for 40 consecutive days. Body weight and epididymal white adipose tissue (EWAT) mass were significantly elevated in HFD and HFD-CO mice compared to LFD mice, but no differences existed between HFD and HFD-CO mice. Area under the insulin-assisted glucose tolerance curve was significantly elevated in HFD and HFD-CO mice compared to LFD mice, but no differences existed between HFD and HFD-CO mice. Heme oxygenase-1 (HO-1) mRNA was significantly higher in EWAT of HFD-CO compared to HFD mice, indicating effective delivery of CO to adipose tissue. CO treatment did not alter *Rev-erb-α* expression or *Rev-erb-α* targets. These results indicate that CO inhalation does not attenuate diet-induced obesity/insulin resistance.

Introduction

Obesity is an increasingly prevalent metabolic disease that affects more than one third of United States adults [1]. Obesity is associated with insulin resistance and increases the risk of developing type 2 diabetes, a disease that results in an estimated \$245 billion in total healthcare costs [2]. Despite the prevalence and health risks associated with obesity, there are few viable and effective treatment options. Therefore, the development of new interventions that can promote weight loss and improve insulin sensitivity are paramount to reducing the prevalence of obesity and its associated diseases.

Recent studies demonstrate that increasing heme oxygenase-1 (*HO-1*) expression can reduce adiposity and heighten insulin sensitivity [3-6]. Although *HO-1* induction by porphyrin molecules or genetic manipulations can improve obesity and insulin resistance, the cellular mechanism responsible for these changes remains to be established. One hypothesis is that the products of HO-1 enzyme activity, CO and biliverdin, exert the metabolically beneficial effects. CO has been shown to reduce inflammatory markers in both cultured cells and mice [7] and a novel CO releasing molecule can attenuate weight gain and preserve insulin sensitivity when mice are placed on a HFD [8]. However, whether or not CO alone can reverse already established obesity and insulin resistance is not established, although a recent study indicates that CO inhalation produces transient reduction in body fat of obese mice [9]. Further, the mechanism by which CO might promote weight loss and improve insulin sensitivity are not known. Establishing the efficacy of CO inhalation to reduce adiposity as well as identifying the mechanism of CO action may lead to novel interventions to combat obesity and type 2 diabetes.

CO may bind to the nuclear receptor *Rev-erb-α* [10]. *Rev-erb-α* is a ligand regulated transcription factor that acts as a transcriptional

repressor [11,12] regulating circadian rhythms and metabolism [13,14]. Although heme has been identified as the ligand for *Rev-erb-α* [15,16], evidence suggests the *Drosophila* orthologue to *Rev-erb-α* binds CO and represses gene expression [10]. Whether or not the expression of *Rev-erb-α* targets is altered in tissues following CO treatment is not known, although a *Rev-erb* agonist has been shown to promote weight loss and suppress PGC-1 α mRNA expression, an established target of *Rev-erb-α* [14].

The purpose of the present study was to determine if CO treatment could reverse obesity and improve insulin resistance due to a high fat diet (HFD) in a manner that was associated with changes in the mRNA expression of well-established *Rev-erb-α* targets in white adipose tissue. We hypothesized that daily CO treatments would reduce adiposity and improve insulin action in mice with established diet-induced obesity. We also hypothesized that mRNA expression of the *Rev-erb-α* targets PGC-1 α , PPAR γ , Bmal1, and *Rev-erb-α* itself would be reduced by CO treatment.

Methods

Treatment of animals

Wildtype C57B6 male mice were purchased from the Jackson

Correspondence to: Thomas H. Reynolds IV, Ph.D., Department of Health and Exercise Sciences, Skidmore College, 815 North Broadway, Saratoga Springs, NY 12866, USA, Tel: (518) 580-8349, Fax: (518) 580-8356; **E-mail:** treynold@skidmore.edu

Key words: insulin, obesity, *rev-erb*, carbon monoxide, diabetes

Received: April 17, 2016; **Accepted:** May 28, 2016; **Published:** May 31, 2016

Laboratory (Bar Harbor, ME). Upon arrival to the animal facility at Skidmore College, all mice (~4-6 weeks old) were housed individually with cage enrichment nest-lets and fed ad libitum chow and water for one week. Mice were then randomly assigned to either a HFD (Test Diets, Catalog #1810251, 60% kcal from fat) or a low fat diet (LFD) (Test Diets, Catalog #58145, 12% kcal from fat). Following 15 weeks of the dietary intervention, HFD-fed mice were body weight-matched-assigned to a CO treatment group (HFD-CO) or control group. Mice from the HFD-CO group were placed in individual induction chambers where they received CO gas (250 ppm) for 60 min per day for 40 consecutive days [17,18]. This CO concentration is slightly higher than what has been shown to increase carboxyhemoglobin levels [9]. Induction chamber CO levels were confirmed by a CO monitor. Control mice from the HFD and LFD groups were placed in the induction chamber for 5 min daily and received air. During the intervention mice were maintained on their HFD or LFD. All animal care and surgery were conducted in accordance with the National Research Council's Guide for Care and Use of Laboratory Animals (Institute of Laboratory Animal Resources, Commission on Life Sciences, 1996). All experimental protocols were approved by Skidmore College's Institutional Animal Care and Use Committee.

In vivo insulin action

To assess the effects of a HFD and CO treatment on *in vivo* insulin action, mice were subjected to an insulin-assisted glucose tolerance (IAGT) test. Prior to the IAGT test mice were fasted overnight. Following the overnight fast, mice were simultaneously administered glucose (2 g/Kg body weight) and insulin (2 U/Kg body weight) via intraperitoneal injection. Glucose was measured by a glucometer (ACCU-CHEK Aviva, Roche Diagnostics) in blood collected via the tail vein at 0, 20, 40, and 60 min following the glucose/insulin injection. We have previously shown that the IAGT test detects insulin resistance as well as an insulin tolerance test in mice fed a HFD, but avoids the severe hypoglycemia that is typically observed during and after insulin tolerance testing [19].

Surgical procedures

Prior to harvesting tissues, mice were fasted overnight and then anesthetized with a 1:1:1 mixture of promace, ketamine hydrochloride, and xylazine by an intraperitoneal injection (0.015 ml/10 g body weight). Epididymal white adipose tissue (EWAT) was rapidly dissected, frozen in liquid nitrogen, and stored at -80° C until analysis.

EWAT RNA extraction and real time quantitative PCR

Total RNA was extracted from EWAT using RNA extraction kits for high lipid content tissue (Qiagen). RNA was quantified by measuring absorbance at 260 nm using a spectrophotometer (Beckman Coulter, Brea, CA). A 1 ug aliquot of total RNA was reverse transcribed using the RETROscript kit from Ambion (Austin, TX). The resultant cDNA (20 ng cDNA/sample in duplicate) was then subjected to quantitative polymerase chain reaction (qPCR) using standard target specific TaqMan gene expression assays for *Rev-erb-alpha* (Assay ID: Mm00441730_m1), HO-1 (Assay ID: Mm00516005_m1), peroxisome proliferator activated receptor gamma (PPARγ) (Assay ID: Mm01184322_m1), PPARγ coactivator-1 alpha (PGC-1α) (Assay ID: Mm01208835_m1), and Bmal1 (Assay ID: Mm00500226_m1) and a real time PCR system (StepOne Plus Real-Time PCR System, Applied Biosystems, Foster City, CA). Relative quantitation of amplified cDNA targets were determined by the ΔΔCT method using StepOne v2.1 software (Applied Biosystems).

Statistical analysis

A one-way analysis of variance (ANOVA) was utilized to detect the effects for diet (LFD vs. HFD) and CO treatment on gene expression. A repeated measures one-way ANOVA was utilized to detect differences in blood glucose levels during the IAGTT. Following a significant F ratio and inspection of interactions, Fisher's LSD post-hoc test was used to locate statistically significant differences between groups. Data are expressed as means ± SEM, and the level of statistical significance was set at p < 0.05. For mRNA expression, means ± SEM were expressed relative to the LFD group mean (% LFD). Data was analyzed using StatView statistical software (SAS Institute Inc., Version 5.0, Cary, NC).

Results

Effect of CO treatment on adiposity

To assess the effects of a HFD and CO treatment on adiposity, we measured body weight and EWAT mass. As expected, a HFD produced a significant increase in body weight and EWAT mass compared to mice fed a LFD (Figures 1A and 1B). Following the 15 week dietary intervention, HFD mice were body weight-match-assigned to a CO treatment group or a control group (room air) and maintained on the HFD. In HFD mice. Treatment with CO (1 hr /day) for did not significantly change body weight or EWAT mass (Figures 1A and 1B), indicating that CO does not reverse pre-existing diet-induced obesity.

Effect of CO treatment on insulin action

In vivo insulin action was determined by conducting IAGT tests




Figure 1. CO treatment does not reduce diet-induced obesity in mice. Body weights were assessed at various time points during CO treatment (A). EWAT mass was assessed at the end of the treatment period (B). *Denotes statistically significant differences from LFD group. N=6-7 mice per group.

that detect diet-induced insulin resistance at more physiologically relevant blood glucose levels than insulin tolerance tests Blood glucose values were higher during the IAGT test in mice fed a HFD compared to a LFD, indicating the presence of diet-induced insulin resistance (Figures 2A and 2B). In HFD mice treated with CO, blood glucose levels were similar during the IAGT test compared to HFD mice treated with room air (Figures 2A and 2B), indicating that CO treatment does not improve insulin resistance in mice with pre-existing diet-induced obesity.

Effect of CO treatment on HO-1 mRNA Expression

Because CO treatment, whether administered by gas inhalation or CO releasing molecules, increases HO-1 expression and activity [8,17,20], we used HO-1 mRNA expression in EWAT as a positive control to demonstrate that our CO gas treatment regimen was effective in adipose tissue. As expected, treating mice with CO gas (250 ppm, 1 hr/day) for six weeks resulted in a significant increase in HO-1 mRNA expression in EWAT from mice fed a HFD (Figure 3).

Effect of CO treatment on the expression of *Rev-erb-alpha* targets

Rev-erb-alpha is a gas responsive transcriptional repressor that has been shown to regulate circadian rhythms, metabolism, and adiposity




Figure 2. CO treatment does not improve insulin-assisted glucose tolerance (IAGT) in mice fed a HFD. Mice were simultaneously injected with 2.0 U/Kg insulin and 2.0 g/Kg glucose and blood glucose values were assessed at baseline and 20, 40, and 60 min following the injection (Panels A) and the area under the IAGT curve was calculated (Panels B). *Denotes statistically significant difference from LFD group. N=6-7 mice per group.




Figure 3. CO treatment increases HO-1 mRNA levels in adipose tissue from mice fed a HFD. The mRNA levels for HO-1 from EWAT of LFD, HFD and HFD-CO mice were quantified by real time PCR as described in Methods. *Denotes statistically significant difference from the LFD group. N=4-5 mice per group.

[10,13,14]. Because CO is thought to bind *Rev-erb* and promote transcriptional repression [10], we examined the mRNA expression of *Rev-erb-alpha* targets in EWAT from mice with pre-existing diet-induced obesity that were treatment with CO for six weeks. CO treatment did not alter the expression of the *Rev-erb-alpha* targets *Bmal1*, *Pparγ*, and *Pgc-1α* (Figures 4A-4C). Since *Rev-erb-alpha* represses its own expression, we assessed its mRNA expression in response to CO treatment and observed no significant changes (Figure 4A). These finding indicate that *in vivo* CO treatment does not alter the expression of established *Rev-erb-alpha* targets in mice with established diet-induced obesity.

Discussion

The present study examined the effect of chronic CO treatment on adiposity, insulin action, and the expression of *Rev-erb-alpha* targets in mice with established diet-induced obesity. We observed no significant changes in adiposity as assessed by body weight and EWAT mass in HFD-fed mice that were treated with CO compared to HFD-fed mice that were treated with air. Further, chronic CO treatments did not improve HFD-induced insulin resistance as blood glucose values during IAGT test were almost identical between HFD-CO and control HFD-fed mice. Finally, the expression of *Rev-erb-alpha* targets were not altered in EWAT of mice that received daily CO treatment. However, CO treatment increased HO-1 mRNA levels in adipose tissue, thereby uncoupling the relationship between HO-1, insulin sensitivity, and weight loss [3].

CO is a product of HO-1 enzymatic activity and thought to mediate the improved insulin action and weight loss observed when the HO-1 gene is overexpressed [3-6]. In this context, the most interesting finding of the present study is the observation that CO treatment increased the expression of HO-1 mRNA without altering insulin action or adiposity. Our present observation is unique and different than studies demonstrating that increasing HO-1 expression by metalloporphyrin administration results in a reduction in body weight [3,21,22]. Importantly, it appears that the ability of metalloporphyrins to decrease adiposity and improve insulin sensitivity is dependent on HO-1 induction because the HO-1 inhibitor, stannous mesoporphyrin, abolished the metalloporphyrin-mediated decrease in body weight [3]. However, other studies support our lack of an effect of CO mediated increases in HO-1 mRNA expression on insulin action and obesity. For example, overexpression of HO-1 in adipose tissue does not alter HFD-induced obesity or insulin resistance in mice [23] and HO-1 expression was a strong predictor of metabolic disease in humans [24]. The present




Figure 4. CO treatment does not alter the expression of *Rev-erb-α* and *Rev-erb-α* targets in adipose tissue from mice fed a HFD. The mRNA levels for *Bmal1* (A), *Pparγ*(B), *Pgc-1α* (C), and *Rev-erb-α* (D) from EWAT of LFD, HFD and HFD-CO mice were quantified by real time PCR as described in Methods. *Denotes statistically significant difference from the LFD group. N=5-7 mice per group.

study provides additional evidence that inducing HO-1 expression, at least by CO inhalation, is not sufficient for improving insulin action or reducing adiposity.

Our results showing that chronic CO inhalation does not improve HFD-induced obesity and insulin resistant may be related to how CO was administered. Hosick *et al.* demonstrated that carbon monoxide releasing molecules promote weight loss and improve insulin action [8]. Although 10 weeks of CO inhalation appears to reduce body weight, this effect is transient and completely absent at 35 weeks [9]. Interestingly, the present results show no effect of approximately six weeks of daily CO inhalation (1 hour/day for 40 days) on body weight or insulin action in mice with established obesity. It is difficult to reconcile the differences between our findings and that of Hosick *et al.* [8,9]. Since some of the beneficial effects of CO are mediated by an increase in reactive oxygen species [25], this response may be abrogated in mice with established weight stable obesity because of the already high levels of ROS [26]. Finally, the lack of a CO effect in the present study and that of Hosick *et al.* [9] may be related to the delivery of the gas by inhalation rather than via CO releasing molecules [8], particularly since CO gas is primarily transported bound to hemoglobin and CO releasing molecules reach tissues independent of hemoglobin. However, we observed a significant increase in HO-1 expression in adipose tissue following the CO inhalation treatment indicating effective CO delivery to cells [8,17,20], although we did not directly measure tissue CO content or assess markers for CO action in other tissues. Future work is needed to determine the mechanism for the different responses between CO delivered by inhalation compared to carbon monoxide releasing molecules.

The present study also tested the novel hypothesis that CO would alter *Rev-erb-α* dependent gene expression. *Rev-erb-α* is a nuclear receptor and thought to be a gas responsive transcription factor that regulates circadian rhythms and metabolism [10,13,14,27]. Recently, a *Rev-erb* agonist has been shown to promote weight loss and suppress PGC-1α mRNA expression, an established target of *Rev-erb-α* [14]. Previous work demonstrated that E75, a *Drosophila* orthologue of human *Rev-erb-α*, binds CO and NO [10]. Subsequent work in cultured HEK 293T and HepG2 cells confirmed that CO and NO bind *Rev-erb-α*, however CO only had modest effects on the expression of *Rev-erb-α* targets compared to NO [27]. The present study clearly demonstrates that CO inhalation does not alter the expression of *Rev-*

erb-α targets in adipose tissue of obese mice. It is unlikely that the gas delivery method explains the ineffectiveness of CO treatment on the expression of *Rev-erb-α* targets because CO inhalation has been shown to stimulate PGC-1α dependent gene expression in skeletal muscle and anti-inflammatory signaling in liver [17,28]. Furthermore, we observed an increase in HO-1 mRNA expression in adipose tissue in HFD-fed mice that were treated with CO. However, it is possible that obese mice are resistant to CO action as we have observed a significant decline in insulin action and PGC-1α in adipose tissue of lean mice that were treated with CO for 7 days (unpublished data).

Conclusion

We have shown that CO treatment does not reverse established diet-induced obesity and insulin resistance despite an increase in HO-1 mRNA levels. Furthermore, CO treatment does not alter the expression of *Rev-erb-α* targets or *Rev-erb-α* itself. Future studies examining the effects of CO releasing molecules on established weight-stable obesity and *Rev-erb-α* are warranted. Because NO binds *Rev-erb-α* with greater affinity than CO, future studies that interrogate the effect of nitrate donors on adiposity and the expression of *Rev-erb-α* targets are needed.

Authorship and contributorship

THR is the principal investigator that designed and directed all aspects of the study. TB, RM, and BS all contributed to the manuscript by conducting experiments, analyzing data, and creating figures. TB and RM assisted in preparing the manuscript for submission.

Funding information

This work was supported by National Institutes of Health 1 R15 DK090722-01, a Skidmore College Opportunity Fund Grant and Skidmore College Summer Research Grant.

Conflict of interest

The authors declare that they have no financial or non-financial competing interests.

References

- Ogden CL, Carroll MD, Flegal KM (2014) Prevalence of obesity in the United States. *JAMA* 312: 189-190. [Crossref]
- American Diabetes Association (2013) Economic costs of diabetes in the U.S. in 2012. *Diabetes Care* 36: 1033-1046. [Crossref]
- Li M, Kim DH, Tsenovoy PL, Peterson SJ, Rezzani R, et al. (2008) Treatment of obese diabetic mice with a heme oxygenase inducer reduces visceral and subcutaneous adiposity, increases adiponectin levels, and improves insulin sensitivity and glucose tolerance. *Diabetes* 57: 1526-1535. [Crossref]
- Burgess A, Li M, Vanella L, Kim DH, Rezzani R, et al. (2010) Adipocyte heme oxygenase-1 induction attenuates metabolic syndrome in both male and female obese mice. *Hypertension* 56: 1124-1130. [Crossref]
- Csongradi E, Docarmo JM, Dubinion JH, Vera T, Stec DE (2012) Chronic HO-1 induction with cobalt protoporphyrin (CoPP) treatment increases oxygen consumption, activity, heat production and lowers body weight in obese melanocortin-4 receptor-deficient mice. *Int J Obes (Lond)* 36: 244-253. [Crossref]
- Cao J, Peterson SJ, Sodhi K, Vanella L, Barbagallo I, et al. (2012) Heme oxygenase gene targeting to adipocytes attenuates adiposity and vascular dysfunction in mice fed a high-fat diet. *Hypertension* 60: 467-475. [Crossref]
- Morse D, Pischke SE, Zhou Z, Davis RJ, Flavell RA, et al. (2003) Suppression of inflammatory cytokine production by carbon monoxide involves the JNK pathway and AP-1. *J Biol Chem* 278: 36993-36998. [Crossref]
- Hosick PA, AlAmodi AA, Storm MV, Gousset MU, Pruett BE, et al. (2014) Chronic carbon monoxide treatment attenuates development of obesity and remodels adipocytes in mice fed a high-fat diet. *Int J Obes (Lond)* 38: 132-139. [Crossref]

9. Hosick PA, Ahmed EK, Gousset MU, Granger JP, Stec DE (2014) Inhalation of carbon monoxide is ineffective as a long-term therapy to reduce obesity in mice fed a high fat diet. *BMC Obes* 1: 6. [[Crossref](#)]
10. Reinking J, Lam MM, Pardee K, Sampson HM, Liu S, et al. (2005) The Drosophila nuclear receptor *e75* contains heme and is gas responsive. *Cell* 122: 195-207. [[Crossref](#)]
11. Harding HP, Lazar MA (1995) The monomer-binding orphan receptor Rev-Erb represses transcription as a dimer on a novel direct repeat. *Mol Cell Biol* 15: 4791-4802. [[Crossref](#)]
12. Zamir I, Harding HP, Atkins GB, Hörlein A, Glass CK, et al. (1996) A nuclear hormone receptor corepressor mediates transcriptional silencing by receptors with distinct repression domains. *Mol Cell Biol* 16: 5458-5465. [[Crossref](#)]
13. Duez H, Staels B (2009) Rev-erb-alpha: an integrator of circadian rhythms and metabolism. *J Appl Physiol (1985)* 107: 1972-1980. [[Crossref](#)]
14. Solt LA, Wang Y, Banerjee S, Hughes T, Kojetin DJ, et al. (2012) Regulation of circadian behaviour and metabolism by synthetic REV-ERB agonists. *Nature* 485: 62-68. [[Crossref](#)]
15. Yin L, Wu N, Curtin JC, Qatanani M, Szewergold NR, et al. (2007) Rev-erbalpha, a heme sensor that coordinates metabolic and circadian pathways. *Science* 318: 1786-1789. [[Crossref](#)]
16. Raghuram S, Stayrook KR, Huang P, Rogers PM, Nosie AK, et al. (2007) Identification of heme as the ligand for the orphan nuclear receptors REV-ERBalpha and REV-ERBbeta. *Nat Struct Mol Biol* 14: 1207-1213. [[Crossref](#)]
17. Beckman JD, Belcher JD, Vineyard JV, Chen C, Nguyen J, et al. (2009) Inhaled carbon monoxide reduces leukocytosis in a murine model of sickle cell disease. *Am J Physiol Heart Circ Physiol* 297: H1243-1253. [[Crossref](#)]
18. Joe Y, Kim SK, Chen Y, Yang JW, Lee JH, et al. (2015) Tristetraprolin mediates anti-inflammatory effects of carbon monoxide on lipopolysaccharide-induced acute lung injury. *Am J Pathol* 185: 2867-2874. [[Crossref](#)]
19. Reynolds TH 4th, Cinquino N, Anthony M, Phelps CB, Zachary Berk E (2009) Insulin resistance without elevated mammalian target of rapamycin complex 1 activity in muscles of mice fed a high-fat diet. *J Appl Physiol (1985)* 107: 1479-1485. [[Crossref](#)]
20. Piantadosi CA, Carraway MS, Babiker A, Suliman HB (2008) Heme oxygenase-1 regulates cardiac mitochondrial biogenesis via Nrf2-mediated transcriptional control of nuclear respiratory factor-1. *Circ Res* 103: 1232-1240. [[Crossref](#)]
21. Galbraith RA, Kappas A (1989) Regulation of food intake and body weight by cobalt porphyrins in animals. *Proc Natl Acad Sci U S A* 86: 7653-7657. [[Crossref](#)]
22. Galbraith RA, Kappas A (1990) Cobalt-protoporphyrin suppresses expression of genetic obesity in homozygous (fa/fa) Zucker rats. *Pharmacology* 41: 292-298. [[Crossref](#)]
23. Huang JY, Chiang MT, Chau LY (2013) Adipose overexpression of heme oxygenase-1 does not protect against high fat diet-induced insulin resistance in mice. *PLoS One* 8: e55369. [[Crossref](#)]
24. Jais A, Einwallner E, Sharif O, Gossens K, Lu TT, et al. (2014) Heme oxygenase-1 drives metaflammation and insulin resistance in mouse and man. *Cell* 158: 25-40. [[Crossref](#)]
25. Zuckerbraun BS, Chin BY, Bilban M, d'Avila JC, Rao J, et al. (2007) Carbon monoxide signals via inhibition of cytochrome c oxidase and generation of mitochondrial reactive oxygen species. *FASEB J* 21: 1099-1106. [[Crossref](#)]
26. Furukawa S, Fujita T, Shimabukuro M, Iwaki M, Yamada Y, et al. (2004) Increased oxidative stress in obesity and its impact on metabolic syndrome. *J Clin Invest* 114: 1752-1761. [[Crossref](#)]
27. Pardee KI, Xu X, Reinking J, Schuetz A, Dong A, et al. (2009) The structural basis of gas-responsive transcription by the human nuclear hormone receptor REV-ERBbeta. *PLoS Biol* 7: e43. [[Crossref](#)]
28. Rhodes MA, Carraway MS, Piantadosi CA, Reynolds CM, Cherry AD, et al. (2009) Carbon monoxide, skeletal muscle oxidative stress, and mitochondrial biogenesis in humans. *Am J Physiol Heart Circ Physiol* 297: H392-399. [[Crossref](#)]